

Copyright WILEY-VCH Verlag GmbH & Co. KGaA, 69469 Weinheim, Germany, 2012.

ADVANCED MATERIALS

Supporting Information

for *Adv. Mater.*, DOI: 10.1002/adma.201204060

Photoalignment and Surface-Relief-Grating Formation are Efficiently Combined in Low-Molecular-Weight Halogen-Bonded Complexes

Arri Priimagi, Marco Saccone, Gabriella Cavallo, Atsushi Shishido,* Tullio Pilati, Pierangelo Metrangolo,* and Giuseppe Resnati**

Photoalignment and surface-relief grating formation are efficiently combined in low-molecular weight halogen-bonded complexes

Arri Priimagi^{1,2*}, Marco Saccone³, Gabriella Cavallo³, Atsushi Shishido^{1*}, Tullio Pilati³, Pierangelo Metrangolo^{3,4,5*}, and Giuseppe Resnati^{3,4*}

Supporting Information

¹ Dr. A. Priimagi, Prof. A. Shishido
Chemical Resources Laboratory, Tokyo Institute of Technology
R1-12 Nagatsuta, Midori-ku, Yokohama 226-8503, Japan
arri.priimagi@aalto.fi; ashishid@res.titech.ac.jp

² Dr. A. Priimagi
Department of Applied Physics, Aalto University of Science and Technology
P.O. Box 13500, FI-00076 Aalto, Finland

³ M. Saccone, Dr. G. Cavallo, Dr. T. Pilati, Prof. P. Metrangolo, Prof. G. Resnati
NFMLab, DCMIC “Giulio Natta”, Politecnico di Milano
Via Mancinelli 7, I-20131 Milano, Italy
pierangelo.metrangolo@polimi.it; giuseppe.resnati@polimi.it

⁴ Prof. P. Metrangolo, Prof. G. Resnati
Center for Nano Science and Technology@Polimi, Istituto Italiano di Tecnologia,
via Pascoli 70/3, I-20133 Milano, Italy

⁵ Prof. P. Metrangolo
VTT, Technical Research Centre of Finland
Tietotie 2, FIN-02044 VTT, Finland

S1. Single crystal X-Ray diffraction analysis.

Crystallographic data of 3. Chemical formula: $C_{28}H_{23}F_4I\text{ON}_4$; $M_r = 634.40$; monoclinic, $P2_1/n$; crystal color: dark-red; shape: wedge; dimensions: $0.12 \times 0.25 \times 0.32$ mm³, $a = 10.2984(15)$, $b = 8.8606(12)$, $c = 28.157(4)$ Å, $\beta = 94.345(12)^\circ$, $Z = 4$, $d_{\text{calc}} = 1.645$ g cm⁻³, $\mu(\text{Mo-K}\alpha) = 1.309$ mm⁻¹, temperature 103(2) K. Crystallographic data have been collected on a Bruker Smart APEX-II CCD diffractometer, equipped with a Bruker KRIOFLEX low temperature device; data collection range $2.06 < \theta < 36.23^\circ$ (completeness = 1.00 for $\theta < 32.00^\circ$), $\theta/2\theta$ scan mode, 97266 collected reflections, 10581 independent, 9057 with $I > 2\sigma(I)$, multi-scan absorption correction: $0.6606 \leq T \leq 0.7471$, $R_{\text{ave}} = 0.0297$. The structure was solved by *SIR2002*^[S1], and refined by *SHELXL*^[S2]. Anisotropic heavy atoms, isotropic hydrogen atoms; H atoms were refined with soft restraints, imposing similar distance on chemically equivalent C-H distances and on H-C-H methyl angles; 412 parameter refined, 178 restraints; final disagreement factors based on all (and ‘observed’) independent reflections $R = 0.0511$ (0.0415) and $wR = 0.0991$ (0.0947), $GOF = 1.134$.




Figure S1: Crystal packing of the supramolecular complex **3** formed upon XB driven self-assembly of **1** and **2**. Ball and stick model. Colors are as follows: C, gray; H, light gray; N, sky blue; F, yellowish green, I, magenta.

S2. X-Ray powder diffraction analysis




Figure S2: Simulated from single crystal data (top) and experimental (bottom) XRPD patterns of **3**

[S1] M.C. Burla, M. Camalli, B. Carrozzini, G.L. Cascarano, C. Giacovazzo, G. Polidori, R. Spagna, *J. Appl. Crystallogr.* **2003**, *36*, 1103.

[S2] G. M. Sheldrick, *Acta Crystallogr. A*, **2008**, *64*, 112.